

# Systems Biology and Bioinformatics in Aging Research: A Workshop Report

Georg Fuellen,<sup>1</sup> Jörn Dengjel,<sup>2</sup> Andreas Hoeflich,<sup>3</sup> Jan Hoeijemakers,<sup>4</sup> Hans A. Kestler,<sup>5</sup>  
Axel Kowald,<sup>6</sup> Steffen Priebe,<sup>7</sup> Dietrich Rebholz-Schuhmann,<sup>8</sup> Bernd Schmeck,<sup>9</sup> Ulf Schmitz,<sup>10</sup>  
Alexandra Stolzing,<sup>11</sup> Jürgen Sühnel,<sup>12</sup> Daniel Wuttke,<sup>13</sup> and Julio Vera<sup>10</sup>

## Abstract

In an “aging society,” health span extension is most important. As in 2010, talks in this series of meetings in Rostock-Warnemünde demonstrated that aging is an apparently very complex process, where computational work is most useful for gaining insights and to find interventions that counter aging and prevent or counteract aging-related diseases. The specific topics of this year’s meeting entitled, “RoSyBA: Rostock Symposium on Systems Biology and Bioinformatics in Ageing Research,” were primarily related to “Cancer and Aging” and also had a focus on work funded by the German Federal Ministry of Education and Research (BMBF). The next meeting in the series, scheduled for September 20–21, 2013, will focus on the use of ontologies for computational research into aging, stem cells, and cancer. Promoting knowledge formalization is also at the core of the set of proposed action items concluding this report.

## Introduction

ON SEPTEMBER 15–17, 2011, a workshop on computational aspects of aging research took place in Rostock-Warnemünde, jointly organized by Georg Fuellen and Julio Vera. It was a sequel to a similar workshop held in 2009.<sup>1</sup> Here we summarize the main results presented at the workshop, including a list of recommendations to foster further progress (see conclusions, below). Figure 1 gives a graphical overview of the connections between the workshop topics, computational analysis methods, some of the molecular entities, and the speakers. The workshop website is [www.ibima.med.uni-rostock.de/IBIMA/symposium/2011/](http://www.ibima.med.uni-rostock.de/IBIMA/symposium/2011/).

Aging research is important in our society, where health-care costs for the aged increase dramatically at a rapid and

highly detrimental pace. Apart from cancer, various other diseases such as dementia and diabetes are closely linked to old age. Just increasing life span by standard medical approaches (which focus on repair, not on slowing down aging by itself) does not necessarily increase health span nor reduce morbidity. On the other hand, influencing the rate of aging just a bit and thereby prolonging average health span would have a huge positive impact for individuals as well as the economy. Because the aging process appears to be so complex, computational approaches are a prudent ingredient in our arsenal of methods toward realizing such an endeavor.

Accordingly, the use of systems biology and bioinformatics to deal with the complexity inherent to aging research has emerged in the last years. Even new research institutes have been created devoted to the idea of using

<sup>1</sup>Institute for Biostatistics and Informatics in Medicine and Ageing Research, Department of Medicine, Rostock University, Germany.

<sup>2</sup>Freiburg Institute for Advanced Studies (FRIAS)–LifeNet, Albert-Ludwigs-University Freiburg, Germany.

<sup>3</sup>Mouse Genetics Group, Leibniz-Institute for Farm Animal Biology (FBN) Dummerstorf, Germany.

<sup>4</sup>Erasmus MC, Dept of Cell Biology and Genetics, Rotterdam, The Netherlands.

<sup>5</sup>Research Group Bioinformatics and Systems Biology, Institute of Neural Information Processing, Ulm University, Germany.

<sup>6</sup>Theoretical Biophysics, Institute for Biology, Humboldt University Berlin, Germany.

<sup>7</sup>Systems Biology and Bioinformatics Group, Leibniz Institute for Natural Product Research and Infection Biology, Hans-Knöll-Institute, Jena, Germany.

<sup>8</sup>European Bioinformatics Institute, EMBL Outstation, Hinxton, United Kingdom.


<sup>9</sup>Chair in Molecular Pulmonology, Forsys Research Group “Systems Biology of Lung Inflammation,” German Center for Lung Research, Philipps-University Marburg, Germany.

<sup>10</sup>Systems Biology and Bioinformatics, Rostock University, Germany.

<sup>11</sup>Fraunhofer Institute for Cell Therapy and Immunology, Leipzig, Germany.

<sup>12</sup>Leibniz Institute for Ageing Research, Fritz Lipmann Institute; Jena Centre for Systems Biology of Ageing-JenAge, Jena, Germany.

<sup>13</sup>Integrative Genomics of Aging Group, University of Liverpool, United Kingdom.



**FIG. 1.** Graphical overview of the RoSyBA 2011 workshop. GWAS, Genome-wide association study; NGS, next-generation sequencing; ROS, reactive oxygen species.

bioinformatics and systems biology to tackle aging. Moreover, in Germany, the federal funding program *GerontoSys* (systems biology for health in old age) currently supports interdisciplinary aging research with a focus on systems biology. *GerontoSys* started with the first call (*GerontoSys1*) at the end of 2009, and in 2011 a second call (*GerontoSys2*) followed. A research program was established that includes three research cores (Cologne, Jena, and Ulm), eight collaborative projects, and three junior research groups. Several of the speakers participating in the workshop are funded by *GerontoSys*; only one collaborative project and one junior research group were not represented.

Aging occurs once organismal growth ceases in the mature animal. The continuous action of the growth program appears to drive aging and with it malignancy (and indeed, cancer risk increases with increasing age). Here, the links between growth, cancer, and aging (normal versus malignant growth/cancer) are apparent. This is also illustrated in the way diet influences organismal growth and aging: Under a low nutrient state, growth is retarded and aging is slowed down. Under conditions of malignant growth, cancer-eliminating mechanisms are important. Fortunately, interventions that extend life span by delaying aging are also often effective against cancer, suggesting that aging itself is intrinsically connected to cancer formation.

By far the strongest prolongation of life span to date has been achieved by influencing the signaling cascades that link the control of growth with nutrient state. The strain specificity of this effect indicates that aging is under strong genetic

control. In mammals, the greatest life span extension so far was achieved by manipulation of a single gene: Mice with a deletion in the *Ghr* (growth hormone receptor) gene lived almost 5 years.<sup>2</sup> Interestingly, in C57BL/6J mice, the same mutation was able to increase life expectancy by only 16%–26%, depending on gender,<sup>3</sup> and the increases amounted to 36%–55% in mice of a mixed genetic background.<sup>4</sup> Thus, it is clear that the genetic environment exerts a tremendous effect on the expression of genetic effects, which may be elucidated by data integration and modeling of genetic effects.

Another motivation for computational approaches is the evolutionary conservation of some of the pathways related to life span control, enabling comparative analyses. An example is the insulin/insulin-like growth factor 1 (IGF-I) signaling (IIS) pathway.<sup>5</sup> A homozygous mutation in the phosphatidylinositol-3-kinase (PI3K) extends life span up to 10-fold<sup>6</sup> in *Caenorhabditis elegans*. In this model, specific roles for the control of life span have been attributed to DAF-2 (ortholog of the insulin/IGF-I receptors), AGE-1 (ortholog of PI3K), and DAF-16 (ortholog of forkhead transcription factors). In *Drosophila melanogaster*, it was shown that INR (ortholog of IGF-I/insulin receptors) or CHICO (ortholog of insulin receptor substrate 1) affect life span as well. Orthologs of components from the IIS pathway have also been described in mammals and in yeast (*Saccharomyces cerevisiae*), where in the latter SCH9 has strong similarities to protein kinase B (AKT) as well as to S6K. The mammalian AKT mediates the effects of PI3K to the forkhead transcription factor family members (FKHR) via phosphorylation.

With respect to the control of life expectancy, regulation of FKHR not only occurs by means of the IGF-PI3K-AKT signaling but also by signals from the mitochondrial or energy metabolic system.<sup>7</sup> Thus, metabolic pathways are critical for patterns of mortality,<sup>8</sup> and caloric restriction (CR) is able to suppress activation of IGF-I, insulin, cytokine, and adiponectin receptors, thus interfering with the onset of cytokine and growth factor-mediated aging processes. Beyond pathways related to metabolism, other aging-associated (sub)networks (e.g., refs. 9–11) are related to telomere length, senescence, and DNA damage response.

### Data Processing and Analysis in the Field of Aging

In the first set of talks, data related to aging were tackled from the viewpoint of bioinformatics and systems biology. Addressing biomedical questions by data integration and subsequent iterative cycles of hypothesis generation and mathematical modeling is a useful approach to analyze high-throughput data, often using interconnected biological networks that are composed by dozens to hundreds of proteins, genes, and microRNAs (miRNAs). Mathematical modeling, rather than being a goal in itself, is a tool to formulate hypotheses, to develop more-directed/better-designed experiments, and to allow predictions. It is useful when dealing with networks enriched in nonlinear motifs like feedback and feed forward loops, which induce nonintuitive regulatory patterns like ultrasensitivity, bistability, and oscillations. The biological networks involved and deregulated in aging are extremely complex and interconnected and display many instances of such nonlinear motifs.<sup>12</sup> As suggested by Kirkwood,<sup>13</sup> there are already a number of examples showing the success of systems biology in unraveling the complexity of aging, in areas such as cell replicative senescence,<sup>10</sup> mitochondrial dysfunction,<sup>14</sup> telomere erosion and DNA damage,<sup>9</sup> and CR.<sup>15</sup>

Although both approaches overlap to a large degree, bioinformatics in our case is considered to include data processing on any level of granularity, based on data sets of any size, whereas systems biology is focused on generating overall (“holistic”) insights from very large data sets on one hand and specific insights by detailed quantitative analyses of selected phenomena on the other hand. Traditionally, bioinformatics finds its roots in computer science, whereas systems biology is closer to mathematical modeling and systems theory.

Large-scale analyses may start with the genome. The genomics of aging was thus reviewed by **Joao Pedro de Magalhaes** (University of Liverpool). Although the human genome was sequenced over 10 years ago, much work remains to decipher it. Unraveling the complexity of the machines of life encoded in the genome is not only essential to understanding biological processes, but is key to manipulating them. In particular, in complex processes such as oncogenesis and aging that involve multiple genes and their interactions with each other and with the environment, a comprehensive understanding is likely to be crucial for developing interventions. Genomic approaches are aimed at increasing our knowledge about how genes and pathways impact on aging and cancer. de Magalhaes presented three examples: (1) The transcriptional profiling of CR to gather insights into its regulation<sup>16</sup> (also see talk by Wuttke, below);

(2) genome-wide approaches using microarray data to identify new candidate genes associated with cancer and aging<sup>17</sup>; and (3) tracking genome evolution to enable an understanding of how different genomes can give rise to species with marked differences in rate of aging.<sup>18</sup>

Genome-wide views of aging gene networks were presented by **Stuart K. Kim** (Stanford University, Stanford). To study aging in mice, he examined expression changes for 8,932 genes in 16 tissues as a function of age. Some tissues displayed large transcriptional differences in old mice, suggesting that these tissues may contribute strongly to organismal decline. Other tissues showed little or no changes in expression with age, indicating strong levels of homeostasis throughout life. He presented a method for the differential analysis of gene co-expression networks and applied this method to look for large-scale transcriptional changes in aging. Comparing the transcriptional profiles for aging in mice to those from humans, flies, and worms, it was found that genes involved in the electron transport chain show common age regulation in all four species, indicating that these genes may be exceptionally good markers of age. However, there is no overall correlation between age regulation in mice and in humans, indicating that the aging process in mice and humans may be fundamentally different. Kim also studies aging in human kidneys. Kidneys age at different rates, such that some people show little or no effects of aging whereas others show rapid functional decline. Whole-genome transcriptional profiling identified 630 genes that change expression with age in the kidney. Expression quantitative trait loci (eQTL) analysis followed by a gene association experiment then identified a single-nucleotide polymorphism (SNP) associated with a hallmark of kidney aging (glomerular filtration rate). The results of this sequential analysis may provide the first evidence for a gene association with kidney aging in humans.

Genome-wide transcriptome analysis with age and the effect of natural genetic variation was the topic of **Ana Viñuela** (Department of Twin Research, King’s College London). To study how genome-wide gene expression regulatory mechanisms progress with age, she first explored genome-wide gene expression variation and regulatory loci (eQTL) in a population of developing and aging *C. elegans* recombinant inbred lines.<sup>19</sup> Throughout the worm’s life, the total number of eQTL decreased with age, whereas the variation in expression increased. Those results suggest that, while for the majority of gene expression becomes more variable during aging, genes that become less variable are more regulated. Moreover, the number of *cis*-acting eQTL in juveniles decreased by almost 50% in old worms whereas the number of *trans*-acting loci decreased by  $\approx 27\%$ , suggesting that aging influences the efficiency of *cis*-regulatory sites for controlling local gene expression. Comparison of a single marker to a multiple marker eQTL model indicated that heritable regulation of gene expression patterns became relatively more complex in old worms due to involvement of multiple loci of small effect.<sup>20</sup> Viñuela is currently studying patterns of gene expression changes and regulation with age in multiple human tissues. Tissue samples were gathered from 855 female Caucasian twins (aged 39–85) recruited from the TwinsUK cohort. The combination of expression profiles with multiple longitudinal age-related phenotypes aims to identify reliable biomarkers for aging and improve

the understanding of gene expression regulation changes with age in humans.

An RNA sequencing (RNA-Seq)-based analysis of aging within a multispecies approach was presented by **Steffen Priebe** (Leibniz Institute for Natural Product Research and Infection Biology, Hans-Knöll-Institute, Jena). This work is part of the JenAge project (see below). Illumina sequencing technology was used for the measurement of the transcriptome. The RNA-Seq data of *Danio rerio* (skin), *C. elegans*, and mouse (brain, liver, and skin) for several time points was mapped versus the reference genomes and an exon-junction database using the Bowtie sequence aligner.<sup>21</sup> Differentially expressed genes (DEG) were identified between each age group and compared across the species. For this reason, orthologous genes were extracted from Ensembl Compara to facilitate the cross-species comparison. Because the individual species exhibit different life spans, the gene expression time courses were rescaled, followed by a combined fuzzy *c*-means clustering of the gene expression profiles to identify common time courses. The optimal number of clusters was estimated by the vote of several cluster validity indices, which capture different aspects of a clustering structure.<sup>22</sup> In this way, gene sets were identified, the expression of which constantly increases or decreases with age in all the three species. Some of them are already known to be associated with aging processes. Functional annotation and pathway analysis were performed to discover functionally related genes. Perturbed gene expression data (rotenone, deoxyglucose) were applied to investigate the influence of mild stress for these genes.

Data and information processing are important aspects of systems biology of aging. **Jürgen Sühnel** from the Jena Centre for Systems biology of Aging–JenAge ([www.jenage.de](http://www.jenage.de)) reported on further JenAge projects in this area. In a first step, the so-called JenAge Information Centre (<http://info-centre.jenage.de/>) was established. It collects and provides general information on aging and age-related diseases as well as on systems biology and is intended to assist researchers in these fields. Although closely linked to the Jena Centre for Systems Biology of Aging, JenAge is a self-contained information resource. It offers a comprehensive compilation of all databases specifically related to aging research, but also collections of books, journals and papers, a meetings calendar, science news, and a list of aging research centers and institutes. In addition, ongoing work includes the setup of a new database, automatic text mining, and data management.

Computational support for systems biology of aging was also provided by **Pat Langley** (Arizona State University). Senescence (see also below) arises from complex interactions among diverse sets of biological processes, and Langley described efforts to encode knowledge about aging in formal computer-based models that allow visualizing causal influences, revising and expanding on these hypotheses, generating explanations for phenomena, and comparing predictions to observations. Users can access, examine, and utilize this content over the internet, as a repository for the community's expanding knowledge about aging.<sup>23</sup>

**Axel Kowald** (Humboldt University Berlin) explained that the BMBF-funded GerontoMitoSys project studies age-related changes in the mitochondria of various model organisms (*S. cerevisiae*, *Podospora anserina*, and *Mus musculus*)

using a combination of experiments and mathematical modeling. One line of research concentrates on the surprising effects seen in *P. anserina* when overexpressing the antioxidant enzyme superoxide dismutase (SOD). Instead of conferring resistance against oxidative stress, the mutants show increased sensitivity and an altered level of antioxidant enzymes (peroxiredoxin) and proteases (CLPP). Incorporating these components into a kinetic model, a positive feedback loop could be identified, demonstrating that the role of superoxide as the primary reactive oxygen species (ROS) responsible for age-related molecular damage is more complicated than originally stated by the free radical theory of aging. This study is a first step toward the integration of the various pathways known to be involved in the control of biological aging.

Modeling the aging epigenome in mathematical terms was the topic of **Jörg Galle** (Interdisciplinary Centre for Bioinformatics, University Leipzig). He presented a computational approach to transcriptional regulation in cells by chromatin-related mechanisms, including histone modification and DNA methylation. In this approach, short-term transcriptional regulation is governed by binding of protein complexes to chromatin, which are capable of reading and writing histone marks. Molecular interactions between these complexes, the DNA, and the histones create a regulatory switch of the transcriptional activity of associated genes that possesses a regulatory memory. Cell proliferation can destabilize this memory and can lead to spontaneous demethylation. Long-term transcriptional regulation is governed by DNA methylation. Considering that methylation of CpGs depends on the modification state of the associated histones, an aging mechanism of the stem cell epigenome is proposed that: (1) Is driven by cell proliferation; (2) induces local hypermethylation of DNA; and (3) affects global transcriptional activity. The proposed model was applied to transcriptional regulation by trithorax complex binding, the recruitment of which has been shown to be CpG density- and methylation state-dependent. This application enables new insights into epigenetic modes of transcriptional regulation. Moreover, it implicates well-founded hypotheses on cooperative histone modifications, proliferation-induced epigenetic changes, and aging of chromatin, all of which await experimental validation.

Boolean networks constitute a qualitative dynamic approach to modeling. **Hans A. Kestler** (Ulm University) spoke about Boolean networks and their evolution for modeling gene regulation. Generating scenarios that model the change of regulatory networks over time can aid in the understanding of the aging process itself. In a typical scenario, an expert models knowledge on genetic interactions as Boolean logic and verifies the resulting hypothetical network by comparing its dynamic behavior to experimental measurements. This is often an iterative process with repeated evaluation and refinement of the network model. Even though the Boolean model is simple, there are still many elements of uncertainty in the modeling process, such as the type of a genetic interaction, or the role of interactions that have not yet been investigated or are hypothesized. Such uncertainties can result in a high number of possible candidate models that have to be investigated by comparing their dynamic behavior with expectations from literature or wet-lab experiments. Supported by the BMBF-funded SyStaR

consortium in Ulm, an approach was developed that supports the generation of aging trajectories by evolving models of gene regulatory networks over time. A preliminary hypothetical network model is specified, and the network evolves according to specified dynamical constraints (*e.g.*, steady states, genes expressed before or after other genes). This is achieved by applying small evolutionary changes to the networks on a symbolic level. Networks are evolved that resemble the input network closely and have desired dynamic properties. The feasibility of our approach was tested on artificially perturbed networks of the mammalian cell cycle and the fission cell cycle. By evolving established network models for early developmental stages, the method can generate models that mimic their age-related evolution.

Metabolic pathway analysis may be considered another approach toward improving our understanding by modeling biological processes. It is applied to nicotinamide adenine dinucleotide (NAD<sup>+</sup>) metabolism with special reference to aging research by **Stefan Schuster** (Department of Bioinformatics, University Jena). NAD<sup>+</sup> is well known as a central cofactor in the redox balance of metabolism. Moreover, NAD<sup>+</sup> is degraded in adenosine diphosphate (ADP)-ribosyl transfer reactions, which are important components in a plethora of signaling reactions. These include reactions linked to DNA repair and aging. Proteins are modified by mono- and poly- (ADP-ribosylation), and histones are subject to NAD<sup>+</sup>-dependent deacetylation catalyzed by sirtuins.<sup>24</sup> Poly-(ADP-ribosylation) is involved in a variety of processes involving DNA transcription, rearrangements, and repair. Several hypotheses on the role of the positive effect of CR on longevity and its relation to NAD<sup>+</sup> metabolism have been put forward.<sup>25</sup> Here, using the concept of elementary flux modes (EFMs)<sup>26</sup> (for a recent review, see ref. 27), Schuster established all potential routes in a network describing NAD<sup>+</sup> biosynthesis and degradation. All known biosynthetic pathways, which include the classical Preiss-Handler pathway and the kynurenine pathway starting from tryptophan, are detected as EFMs. Moreover, several elementary modes are found that degrade NAD<sup>+</sup>, represent futile cycles, or have other functionalities. The systemic and systematic analysis and comparison of the networks specific for yeast and humans document significant differences between species with regard to both the use of precursors, biosynthetic routes, and NAD<sup>+</sup>-dependent signaling.<sup>28</sup> This enables critical examination of the hypothesis that CR increases NAD<sup>+</sup> turnover without altering steady-state NAD<sup>+</sup> levels.

**Niels Grabe** (University of Heidelberg) described a method for the automatic inference of causal networks and the response to growth factors. After automatically integrating the comprehensive set of all known cellular networks from the PID Pathway Interaction Database, off-switched genes are identified and removed, and a subnetwork with the given set of growth factors as the input layer and the desired set of functional cell states as the output layer is extracted. Highly specific networks for relevant signaling cascades result from further restricting the network by differentially expressed genes obtained from stimulation experiments with the growth factors of the input layer. A reverse version of this algorithm is designed to identify those receptors and cell-cell communication pathways altered by aged skin fibroblasts.

Finding cross-links and mechanistic links in biological networks is the bioinformatics task handled by **Bianca Habermann** (MPI-AGE Cologne) to aid biologists in analyzing and interpreting their data. With the advancement of genome- and proteome-scale techniques, this task becomes more and more challenging. Eventually, a genome-wide screen should not exist by itself, but should be consolidated with the vast amount of genome-scale data that are freely available. As a member of the BMBF-funded SyBACol project, Habermann is involved in the integration of expression studies in different model systems. One of her aims is to identify crosstalk between molecules and pathways in biological interaction networks. Finding the most prevalent path between two proteins that are mechanistically linked and the most likely cross-linkers between two functionally active pathway modules is one aim. Another aim is to consolidate data from different species and invest in the functional annotation of hits from expression screens. Although many genetic and high-throughput functional data are only available for model organisms like worm or fly, some players will be equally important in vertebrates or mammals, including humans.

**Jörn Dengjel** (University of Freiburg) outlined that human aging may be studied by specifically investigating organ aging, the kidney being a prime target of age-associated organ damage as reflected by the clear association of renal function decline with age. The BMBF-funded NephAge consortium in Freiburg hypothesizes that the aging process of the kidney is determined by the life span of podocytes, which are terminally differentiated, highly specialized kidney cells. To unravel the molecular identity of kidney aging and responsible signaling networks, NephAge performs a systems biology approach using complementary mouse models, among them the first available model of kidney aging, to isolate primary podocytes for an in-depth proteomic, genomic, and metabolomic characterization. In a first bioinformatics step, the data sets are compared to recognize parallel regulations and patterns and to rapidly deduce first connections between different aging states. Next to the unbiased approaches using mouse models, genes associated with the aging kidney, identified from genome-wide association studies in a large number of middle-aged and elderly individuals, will be studied in genetically tractable and screenable fly and worm models for their role with respect to renal function, autophagy, and longevity. To translate all generated data into predictive dynamic models, NephAge employs complexity reduction strategies to identify and model key signaling pathways crucial for kidney aging. The goal is to identify signaling subsystems and establish a hierarchy of time-graded responses that determine short-term age-related function and long-term age progression.

An emerging topic in aging research are miRNAs, as well as integrating bioinformatics and systems biology to elucidate the role of miRNAs in cancer and aging, which are studied by **Ulf Schmitz** (Department of Systems Biology and Bioinformatics, University of Rostock). miRNAs are a class of small noncoding RNAs that can posttranscriptionally regulate the expression of many protein-coding genes. Once processed, the mature  $\approx 22$ -nucleotide-long molecule represses its mRNA target in conjunction with the RNA-induced silencing complex (RISC) by either inhibiting translation or by mediating mRNA decay. miRNA and target

deregulation has been shown to play a causal role in many aging-related diseases, the one most extensively investigated being cancer.<sup>29</sup> It has been suggested that miRNAs operate at different levels of aging.<sup>30</sup> This study claims that miRNAs are regulated during cellular senescence *in vitro* and contribute to tissue regeneration by regulation of stem cell function; several miRNAs modulate life span in *C. elegans*. In his talk, Schmitz discussed how the integration of bioinformatics and systems biology tools is required to tackle complex biochemical networks involving the regulation by multiple miRNAs. He used the case study of miRNA repression of the cell cycle regulator p21, which is an important player in senescence.<sup>31</sup>

### Growth, Cancer, Inflammation, and Aging

The second set of talks demonstrated how aging is connected to growth, in itself a fundamental process of life, and to its abnormal variant, that is, cancer (see also the Introduction section). Furthermore, all three processes are modulated by inflammation. Comparing these processes on the molecular level by computational means may reveal a set of molecular species (such as proteins) that are jointly involved (see also Fig. 1). Such “anchoring points” may include the somatotrophic axis—IGF-1, PTEN,<sup>32</sup> and mammalian target of rapamycin (mTOR).<sup>33,34</sup>

Genome maintenance and its impact on aging and disease were highlighted by **Jan Hoeijmakers** (Erasmus MC, Rotterdam). An intricate genome maintenance machinery has evolved to counteract the consequences of DNA damage, affecting the function of our genes. Nucleotide excision repair (NER) is one of the most versatile repair systems removing a wide range of helix-distorting lesions, mostly from exogenous sources (ultraviolet [UV], bulky adducts), but also of endogenous origin (oxidative damage, *e.g.*, cyclopurines). It does so in a multistep “cut-and-patch”-type of reaction involving more than 30 proteins. Two NER subpathways exist. Global genome NER operates genome-wide removing lesions, which otherwise would give rise to mutations during replication and hence is critical for preventing mutations. Transcription-coupled repair removes damage that obstructs transcription, primarily counteracting cytotoxic effects of DNA injury.

UV-sensitive inherited NER syndromes display a striking clinical heterogeneity, including very strong (skin) cancer predisposition in xeroderma pigmentosum (XP) and dramatic neurodevelopmental deficits such as in Cockayne syndrome (CS) and trichothiodystrophy (TTD). Different mutations in NER helicases XPB and XPD, subunits of the repair/transcription factor TFIIH, give rise to all three disorders or combinations. Detailed analysis of XPD<sup>TTD</sup> mice, carrying a point mutation of a TTD patient, revealed that TTD (and CS) are segmental premature aging syndromes, with reduced cancer susceptibility. XPD<sup>XP/CS</sup> mutant mice on the other hand are highly predisposed to cancer and also display premature aging, demonstrating that both phenotypes can co-exist. Other single and double NER mutants exhibit multiple premature aging features, including *bona fide* osteoporosis, dramatic neurodegeneration, early infertility and cessation of growth, liver and kidney aging, deafness, retinal photoreceptor loss, exhaustion of hematopoietic stem cells, *etc.* Life span is limited to 1.5 years for milder mutants and to 3–5 weeks for dramatic (double) mutants.

A striking correlation is found between severity and type of defective repair and rate of onset, severity, and type of aging symptoms, providing strong experimental support for the DNA damage theory of aging. Conditional tissue-specific repair mutants target accelerated aging to specific organs, *e.g.*, dramatic neurodegeneration occurring only in the cerebellum, revealing organ-specific accelerated aging. Hoeijmakers proposed that endogenous DNA injury hampers transcription/replication and triggers cellular apoptosis-senescence causing aging. Microarray and functional and physiological studies have revealed that persisting DNA damage elicits a systemic downregulation of the IGF-I somatotrophic axis and upregulation of antioxidant defenses, favoring maintenance and defenses at the expense of growth and development, explaining the severe growth defect of repair mutants. Persisting DNA damage triggers this “survival” response in a cell autonomous manner and implicates regulation by miRNAs. CR and fasting trigger a similar “survival” response, which maximizes antioxidant defense and, when constitutive, promotes longevity, at least under laboratory conditions. These data link accumulation of DNA damage and the IGF-I control of life span and open perspectives for the promotion of healthy aging.

Interactions among growth, metabolism, and aging are also investigated by **Andreas Hoeflich** (Laboratory for Mouse Genetics, Leibniz-Institute for Farm Animal Biology Dummerstorf, Germany). In mouse knockout models lacking Ghr or insulin-like growth factor I receptor (Igf-IR), prominent increases of life span have been described and, thereby, a negative correlation of growth and aging was established.<sup>2,35</sup> However, the lack of one or both receptors is an extremely rare condition in mammals and is not a candidate for acute or adaptive life span elongation *in vivo*. Thus, Hoeflich et al. have started to characterize interaction of growth and life span in phenotype-selected mouse lines characterized by extreme growth, confirming the intricate negative relationship between growth and life span in non-inbred strains (Dummerstorf phenotype-selected mouse strains), a highly interesting set of mouse models useful for the analysis of growth and aging. Whereas both growth and aging represent central physiological features, the exact mechanisms linking both phenomena are still not well understood. As discussed in the Introduction, a central age-related pathway evolutionarily conserved from worms to mammals is the IIS pathway. Because IIS includes a series of receptor tyrosine kinases, including not only the IGF-IR but also the insulin receptor, focus is on the activation of this central signaling pathway.

A longitudinal study was performed in Dummerstorf DU6P mice characterized by extreme muscle mass, selected over 138 generations for high carcass protein amount. In muscle from 29-week female DU6P mice, a massive activation of AKT compared to controls was found. Potential mechanisms of AKT activation include IGF-I, IGF-II, insulin/Igf-receptor, myostatin, or integrin-linked kinase. However, only PTEN, which was severely downregulated in 29-week-old female DU6P mice, could be identified as a potential trigger, consistent with higher levels of AKT-Ser473 phosphorylation.<sup>36</sup> AKT activation also translated into specific inactivation of glycogen synthase kinase 3 $\beta$  and a seven-fold increase of muscular glycogen (5% of total muscle weight). The mouse model described thus comprises phenotypes of

massively elevated growth, severely reduced life span (less than 78% versus controls), and, last but not least, significant metabolic abnormalities (higher protein and glycogen content in muscles), which in total may be related to the altered muscle AKT activity. In 29-week-old female DU6P mice, PTEN might serve as a superordinated pacemaker of growth, aging, and metabolism.

Chronic inflammation has been linked to cancer as well as aging. **Bernd Schmeck** (University of Marburg) demonstrated its relevance in diseases such as diabetes, atherosclerosis, inflammatory bowel diseases, and lung diseases. Among the 10 most important causes of death worldwide are four lung diseases with inflammatory aspects—pneumonia, obstructive lung diseases, tuberculosis, and lung cancer. Obstructive lung diseases, like chronic bronchitis and bronchial asthma, can be caused exogenously by cigarette smoke, air pollution, and allergic agents, as well as bacteria and viruses. Most importantly, this inflammation is not limited to the lung, but a “systemic spillover” seems to affect many organ systems, causing “aging-related” phenotypes like dementia, chronic heart diseases, muscle weakness, and osteoporosis. The aim of the Marburg group is to understand physiological and pathological regulatory circuits of lung inflammation at the crossroads of host defense and tissue destruction, as well as proliferation and aging. Therefore, the molecular machineries of chromatin modifications and small non-protein-coding RNAs/miRNAs are studied, combining wet lab experiments, high-throughput facilities, and bioinformatics, as well as mathematical modeling, to construct a predictive model that may help to identify crucial signaling hubs as possible therapeutic targets for these widespread pulmonary diseases causing preterm aging.

### Influence of Diet on Aging

Manipulating aging and, at the same time, preventing cancer growth and inflammation may best be done by nutritional interventions. Many of the genes discussed above are in fact part of the signaling cascade of nutrient-sensing systems such as growth hormone (GHR) and insulin-like signaling (IGF1, PI3K, PTEN, FoxO) as well as mTOR. The computational study of such interventions can pave the way to understanding the influence of diet on the aging process and thus enable the development of interventions tweaking this system, to interfere with life span via conserved mechanisms active in organisms from yeast to mammals.

Comparative interactomics of dietary restriction (DR)-essential genes is done by **Daniel A. Wuttke** (Integrative Genomics of Ageing Group, University of Liverpool). DR, limiting certain factors in diet without causing malnutrition (also often called CR), is the most powerful nongenetic intervention that delays the aging process and extends healthy life span in multiple organisms from yeast to rodents. To decipher the mechanisms of DR, Wuttke established a web-accessible database (GenDR) of DR-essential genes, which if genetically altered interfere with the effect of DR to extend life span in model organisms (yeast, worm, fly, and mice). Orthologs in the respective other organisms were identified and their molecular evolution investigated, showing that DR-essential genes have more orthologs and in mammals were subject to stronger purifying selection than expected by chance. Molecular interactions with DR-essential genes/

proteins and their orthologs were retrieved from available databases and networks constructed. Basic topology analysis revealed that DR-essential genes tend to be located in the center of the interactome and have a higher interconnectivity to each other than expected by chance. Genes interacting significantly with DR-essential orthologous groups between multiple species, and novel DR-essential genes, which were not previously implicated in the mechanisms of DR-mediated life span extension, were identified via guilt-by-association and experimentally validated. This revealed new insights into the fundamental mechanisms in which the nutritional state impacts on life span determination.<sup>37</sup> GenDR can be found at <http://genomics.senescence.info/diet/>.

Moving from genes to interventions, mining genome-wide drug response compendia to discover novel CR mimetics is done by **Kristen Fortney** (Department of Medical Biophysics, University of Toronto, Toronto, Canada).<sup>38</sup> Genome-scale data on the responses of human cell lines to over 1,000 drug treatments have become available, and their integration with gene expression signatures of CR, pathway information, and protein–protein and drug–target interaction networks yield a prioritized list of candidate CR mimetics. These candidates are annotated using drug–target, pathway, and protein interaction databases, and clustering characterizes them in terms of common chemical properties and of common target proteins, pathways, and diseases. Modes of action can then be compared with those of known life span-affecting drugs, to rank the candidates in terms of their therapeutic promise.

### Stress Response, Senescence, ROS, and Mitochondria

Up to now, growth, cancer, and inflammation were discussed as the fundamental processes related to aging, but stress response should not be forgotten, because it can lead to both life span extension or senescence and subsequent aging. Cellular stress and in particular the “oxidative stress,” caused by ROS, is maybe of highest relevance. ROS and the damage it induces may cause stress responses. While the damage may be detrimental, ROS “signaling” and stress response may also include healthy and longevity-promoting aspects. Senescence is an important aging-related cell phenotype that might result from (oxidative) stress, and mitochondria play an important role here. Senescence may be considered the counterpart of growth, and both may be equally detrimental to health and longevity.

In particular, the connections between mitochondrial dysfunction and cell senescence were highlighted by **Thomas von Zglinicki** (Centre for Integrated Systems Biology of Aging and Nutrition, Newcastle University). Cell senescence is a permanent, irreversible proliferation arrest resulting from, *e.g.*, telomere shortening and persistent DNA damage signaling. In addition to growth arrest, senescent cells display major changes in multiple phenotypes, such as gene expression patterns, metabolism, morphology, and function, which can significantly spread their impact in tissues. Mitochondrial dysfunction (increased ROS production despite decreased mitochondrial membrane potential) is an early hallmark of senescence. Combining probabilistic functional interactome network analyses and serial knockdowns of candidate genes, a signaling pathway was identified, linking DNA damage response to mitochondrial ROS production.

Stochastic modeling suggested that this pathway would be necessary and sufficient for maintenance of proliferation arrest; this was confirmed experimentally for the first developmental phase of senescence. Studies in mice have shown increasing fractions of cells bearing senescence markers in various tissues during aging. Dietary restriction reduced frequencies of senescent cells in the same tissues.<sup>39</sup>

Mitochondrial genome and aging are investigated by a systems biology approach in the BMBF ROSage project presented by **Saleh Ibrahim** and **Georg Fuellen** (University of Lübeck Medical School/University of Rostock Medical School), postulating that: (1) Aging of an organism depends on specific aging-related changes in specific organs, which in turn depend, in part, on specifics of the electron transport chain (ETC) and (2) ROS mediate some aspects of this dependence. To study these hypotheses, mouse strains with well-defined and stable mutations in the ETC yield standardized readouts for several organs, following the mouse strains over time. In parallel, multiple approaches are pursued to model the underlying processes mathematically on the microscale. Bioinformatics data integration is designed to provide connections between the underlying processes and to some of the readouts, and to perform overall machine learning on the whole dataset. The latter will be based on readouts as well as modeling results, using ontologies, towards a better understanding of aging on the macroscale.

A complementary approach is the BMBF-funded project OXISYS, presented by **Katrin Zeilinger** (Berlin Brandenburg Center for Regenerative Therapies [BCRT], Charité Universitätsmedizin Berlin), investigating the role of oxidative injury in aging and therapeutic implications. There is increasing evidence that oxidative stress plays a major role in aging-related processes in the human organism. Within OXISYS, the effect of aging and the role of ROS exposition in aging processes using the liver as a model organ are investigated. The project is conducted in a consortium of four academic partners (Charité Universitätsmedizin Berlin, Eberhard-Karls Universität Tübingen, Universität des Saarlandes, Hans-Knöll-Institut Jena) and two industrial partners (Pharmacelsus GmbH, Insilico Biotechnology AG). The *in vitro* approach is based on three-dimensional perfusion culture of liver cells in a hollow fiber capillary membrane bioreactor providing decentralized mass exchange and integral oxygenation. The system serves as a physiological model for human-predictive studies on the effect of hepatic aging and oxidative stress effects on hepatic metabolism and the contribution of different cell populations to these processes. Rats of different age classes are used to analyze the *in vivo* response to oxidative stress. Biochemical, metabolic, and gene expression data generated in the experimental models will serve for developing *in silico* models to describe, simulate, and predict aging processes in the liver. Models generated from rat *in vivo* and *in vitro* models will be translated to the human liver and verified for their systemic relevance. The models established will be used to identify potential biomarkers and targets for diagnosis and therapeutic regulation of ROS induced tissue aging and degeneration.

Computational advances in the study of senescence, in particular on the role of lipid rafts in aging and cellular senescence, were reported by **Fiete Haack**, (Faculty of Computer Science, University of Rostock).<sup>40</sup> The plasma

membrane of eukaryotic cells is a highly compartmentalized structure, characterized by large, stable, multireceptor complexes, actin-mediated cytoskeleton fences, and mobile lipid rafts. In recent years, lipid rafts in particular have drawn much attention due to their ability of selectively concentrating proteins and thereby facilitating the assembly of signaling complexes. Recent studies indicate an involvement of lipid rafts in cellular senescence and age-associated dysfunction of, *e.g.*, T cell receptor signaling. Unfortunately, biochemical approaches to study lipid rafts in biological membranes are highly limited. Standard techniques, like detergent-resistant membrane or cholesterol-depletion experiments have too complex side effects to draw substantive conclusions. Therefore, computational modeling is an attractive alternative. However, the ascribed static and dynamic organization of plasma membranes leads to highly inhomogeneous spatial distribution and demands a thorough consideration of space when modeling and simulating such systems. Haack and colleagues propose a new modeling and simulation approach that allows integration of different types of spatial dynamics, *e.g.*, compartmental dynamics, mesh-based approaches, or moving individuals, within one model. The associated simulator combines Gillespie, the Next Subvolume method, and Brownian dynamics. This allows the simulation of complex spatial dynamics like those involved in studying the dynamics of lipid rafts and their role in receptor co-localization.

### Stem Cell Aging

With respect to cancer, growth, inflammation, stress response, and aging, the cells with the most prominent role may be various types of stem cells. Stem cell differentiation is tightly controlled by ROS signaling, and stem cells have been hypothesized to be crucial in both maintaining youthful proliferation and regenerative capacity as well as (when out-of-control) to be a driving force of malignancy because of their undifferentiated state.

For example, aging processes in neural stem cells that are affected by the pool of adult stem cells were discussed by **Anne Brunet** (Department of Genetics, Stanford University). In the nervous system, neural stem cells are thought to be critical for learning and memory. During aging, both the pool of neural stem cells and their ability to give rise to new neurons decline, raising the possibility that neural stem cell depletion may underlie part of the cognitive dysfunctions during aging. However, the mechanisms that regulate the neural stem cell pool during aging are largely unknown. Brunet's hypothesis is that genes that regulate life span maintain the homeostasis of adult stem cell pools in long-lived species. Transcription factors of the FoxO family play a conserved role in controlling longevity downstream of the insulin pathway. FoxO orthologs are already known to extend life span in invertebrates and SNPs in the *FoxO3* gene are associated with extreme longevity in humans. Brunet's group has recently found that the transcription factor FoxO3 maintains the neural stem cell pool in adult mice. Specifically, genome-wide analysis of FoxO3 binding sites in neural stem cells using next-generation sequencing technologies identified a network of FoxO3 targets involved in "stemness" and aging. This global analysis has also provided key insight into novel co-factors and chromatin marks that interact with



FoxO3 to regulate adult neural stem cell function. FoxO3's ability to control an epigenetic network that is pivotal for adult neural stem cell homeostasis might help counter brain aging in long-lived species, including humans.

Homeodynamic fitness and the age-related disbalance in mesenchymal stem cell renewal and progenitor molding was the topic of **Günther Lepperdinger** (Institute for Biomedical Aging Research, Innsbruck). Aging of somatic tissues comes along with a decline of regenerative capacity. Tissue regeneration and repair involve the consecutive emergence and parallel integration of new parenchymal cells, which descend from undifferentiated precursors. Multipotent stromal progenitor cells, also known as mesenchymal stem cells (MSCs), are pertinent tissue-specific stem cells in adults. The concept of MSCs is interesting because this particular type of precursor can bring forth a large spectrum of cell types as diverse as bone, cartilage, tendon, or fat precursor cells. Besides these descendants, interstitial stromal cells and fibroblasts, which are also terminally differentiated cells common and widespread throughout the body, are reckoned to be direct offspring of MSCs. These particular stem cells are considered highly effective therapeutic assets to combat a wide spectrum of diseases, and thus approaches with the aim of rebuilding damaged or diseased organs and body parts has yielded a virtually endless list of only recently initiated clinical trials. Assuming a fit cell population can withstand accrued damage and compounding of molecular changes longer than irreversibly aged cells, the interest is to determine means and measures for investigating molecular mechanisms underlying this concept, and in particular to elucidate endogenously expressed bioactive factors and their mode of actions which enhance the fitness of tissue-borne MSCs.

In the discussion of MSCs aging, we are faced with two interrelated aspects, the effect aging has on MSCs themselves and the contribution MSCs have to the aging of the organisms. To complicate the matter, to the first aspect we also have to add the role of the niche in MSC aging. **Alexandra Stolzing** (Fraunhofer Institute for Cell Therapy and Immunology, Leipzig) spoke about the changes in the signal pathways in MSCs isolated from adipose tissue of donors ranging from 20 to 80 years of age. Adipose tissue has been shown to contain high numbers of MSCs (adipose-derived stem cells; ADSCs), which are an alternative source of MSCs for therapeutic applications. ADSCs have similar characteristics compared to bone marrow-derived MSCs (BM-MSCs) and might age in a similar fashion. One of the signaling pathways regulating MSC differentiation, pluripotency, and migration is the WNT signaling pathway. ADSCs were isolated from liposuction and subcutaneous tissue from donors of different age and gender. A dramatic decrease in WNT gene expression has been detected for ADSC-derived from females at the age of 40–60 years. Gene expression analysis from ADSCs derived from liposuction resulted in absence of nearly all WNT genes analyzed and different patterns of cell cycle genes and differentiation markers compared to subcutaneous ADSCs. Thus, the source of the ADSCs has a great impact on cell behavior and therefore for therapeutic applications. Further study of these differences may lead to important discoveries regarding the mechanisms of WNT signaling in cells from different niches and the regeneration capacity of MSC.

## Conclusions

In the future, we believe that aging research will benefit from ever more synergies between experimental work, bioinformatics, and systems biology approaches. Bioinformatics provides a selection of tools, databases, and approaches ideal for compiling existing information regarding the molecular components of aging-related biological networks and their association with pathological conditions (*e.g.*, putative and validated protein–protein and miRNA–mRNA interactions, transcriptional regulation, genetic interactions, *etc.*). Bioinformatics and genomics also offer the tools to obtain, via high-throughput data analysis, empirical evidence suggesting connections between the components of these aging-related networks. The systems biology view complements this approach by making use of this information and organizing it into detailed maps of the regulatory networks investigated. Those maps, when translated into mathematical models, become excellent tools for integrating high-throughput data of highly interconnected biological networks and testing the consistency of new data against the state of the art. But these models are also useful for formulating mechanistic hypotheses on putative interactions explaining inferred links between genes and proteins, and thereby for developing more directed and better-designed experiments for testing them. Furthermore, mathematical modeling is a necessary tool when dealing with networks enriched in nonlinear motifs like feedback and feed forward loops. These regulatory motifs, which are commonly found in the regulatory networks involved in aging and aging-associated diseases, induce nonintuitive regulatory patterns like ultrasensitivity or bistability, impossible to detect and understand without modeling. Their deregulation threatens or alters the natural homeostasis in tissue regeneration, which is necessary to keep functionality and viability and avoid emergence of aging-related phenotypes.

Therefore, we predict a future of aging research in which bioinformatics is used to inform and to construct better, more precise networks that explain in detail the most relevant processes underlying aging conditions. Both, bioinformatics and mathematical modeling are then jointly used to determine the regulatory chain of causation underlying the phenotypic and genotypic patterns obtained from the high-throughput data. A faster and more reliable interpretation of biological data and evidence will be a desired outcome of this synergy. Thus, we are convinced that the two communities will highly benefit from sequels to meetings such as RoSyBA–Rostock Symposium on Systems Biology and Bioinformatics in Ageing Research, in which the most challenging bioinformatics and systems biology tasks are tackled through fruitful and productive alliances between both communities.

Furthermore, we perceive that both communities would highly profit from synergies induced by integrative efforts that result from formalizing the biomedical domain knowledge—biomedical ontological resources are an essential contribution toward this aim.<sup>41</sup> Core ontologies include the phenotypic representation and the annotation of biological processes (such as diseases) with phenotypic features, supported by automated literature analysis. Machine-based reasoning can then guide data integration, consistency analysis, and inference of novel findings, such as relevant

phenotypes as well as diagnostic and interventional suggestions. The cross-species ontological integration of phenotypes is the next key step to achieving data interoperability across resources, inference of core findings, and translational medicine. These results could be exploited for intervention (drug) development across model species and for the reduction of animal experiments. The application of findings from model organisms to humans requires relevance testing and also risk assessment, which could be done with the same framework. Furthermore, it is of high importance to use phenotypic descriptions to formalize the evidence of underlying mechanisms for models describing the development of biological (age/disease) status. Again, this can be based on the conceptual framework for phenotypes. The causal link between the phenotype at different conceptual levels and its genetic predisposition will then be the result of “causal chains,” where one feature, such as the molecular function of a gene, is transposed to a tissue and the organ, leading to the anatomical function for the macroscopic phenotype. Considering the importance of ontologies for progress in computational aging research, the organizers of the 2011 workshop are planning a focus on ontologies for the next workshop in 2013.

Finally, we recommend the following action items:

1. Because aging processes affect all aspects of life, computational analyses require integration of a maximum amount and variety of reliable and accurate biological data and knowledge, and of the integration. Thus, it is proposed to build a (wiki-based) collaborative resource that provides such data integration on all levels. Preparation for this resource must start today, striving for interoperability and open standards. Data deposit will be encouraged by making it mandatory for publication (following an open data paradigm). Validity of the integration will be enhanced by using detailed ontologies. Aging-related computational research must then point out those processes of life that are most fundamental for aging and create computational models for them.
2. Funding must be made possible at sustained levels, beyond Germany and the European Union. Aging research toward health span extension shall be designated as an area of research that has high returns on investment, and computational support is an essential part of it. Molecular aging research shall also be advertised as one “prime application area” for bioinformatics and systems biology data integration efforts in general, to the European Bioinformatics Institute (EBI), the National Center for Biotechnology Information (NCBI), *etc.*
3. Funding for aging research in general should focus more on the impact of early life interventions on life span, and on aged animal cohorts covering an array of species, including non-inbred models. Following these two recommendations would increase the practical relevance of both experimental and computational aging research.

### Acknowledgments

The workshop was kindly sponsored by the German Federal Ministry of Education and Research (BMBF, project CALSYS-FORSYS FKZ. 0315264), the SENS Foundation,

Sanofi-Aventis, and the Department of Medicine and the Department of Aging Science and Humanities (Interdisziplinäre Fakultät), University of Rostock. This work was also supported by the German Federal Ministry of Education and Research (BMBF) within the framework of GerontoSys (Forschungskern SyStaR, project ID 0315894A) to H.A.K., by the Kooperationsprojekt GerontoMitoSys, FKZ 0315584 to A.K., by the Verbundprojekt ROSAge, FKZ 0315892A to G.F., and within the framework FORSYS Partner (Forsys Lung, FKZ 0315256) to B.S. We thank Mark Möller for preparing a first version of the figure.

### Author Disclosure Statement

No competing financial interests exist.

A.S., A.K., B.S., A.H., J.H., H.A.K., J.D., and J.S. contributed on those sections that include the topic of their talks. J.V. and U.S. contributed to Systems Biology of Aging and Conclusions sections. S.P. also contributed to the Introduction and Abstract. D.R.S. contributed to the Conclusions. D.W. worked on Abstract, Introduction, and section introductions. G.F. assembled the report, and contributed to the Abstract, Introduction, section introductions, and Conclusions.

### References

1. Fuellen G, Adjaye J, de Grey A, Lepperdinger G, de Magalhaes JP, Suhnel J, Wipat A. Bioinformatics in aging research: A workshop report. *Rejuvenation Res* 2010;13:763–767.
2. List EO, Sackmann-Sala L, Berryman DE, Funk K, Kelder B, Gosney ES, Okada S, Ding J, Cruz-Topete D, Kopchick JJ. Endocrine parameters and phenotypes of the growth hormone receptor gene disrupted (GHR<sup>-/-</sup>) mouse. *Endocr Rev* 2011;32:356–386.
3. Coschigano KT, Holland AN, Riders ME, List EO, Flyvbjerg A, Kopchick JJ. Deletion, but not antagonism, of the mouse growth hormone receptor results in severely decreased body weights, insulin, and insulin-like growth factor I levels and increased life span. *Endocrinology* 2003;144:3799–3810.
4. Zhou Y, Xu BC, Maheshwari HG, He L, Reed M, Lozykowski M, Okada S, Cataldo L, Coschigano K, Wagner TE, Baumann G, Kopchick JJ. A mammalian model for Laron syndrome produced by targeted disruption of the mouse growth hormone receptor/binding protein gene (the Laron mouse). *Proc Natl Acad Sci USA* 1997;94:13215–13220.
5. Hamet P, Tremblay J. Genes of aging. *Metabolism* 2003;52(Suppl 2):5–9.
6. Ayyadevara S, Tazearslan C, Bharill P, Alla R, Siegel E, Shmookler Reis RJ. Caenorhabditis elegans PI3K mutants reveal novel genes underlying exceptional stress resistance and lifespan. *Aging Cell* 2009;8:706–725.
7. Vijg J, Campisi J. Puzzles, promises and a cure for ageing. *Nature* 2008;454:1065–1071.
8. Barzilai N, Huffman DM, Muzumdar RH, Bartke A. The critical role of metabolic pathways in aging. *Diabetes* 2012;61:1315–1322.
9. Shachar R, Ungar L, Kupiec M, Ruppin E, Sharan R. A systems-level approach to mapping the telomere length maintenance gene circuitry. *Mol Syst Biol* 2008;4:172.
10. Passos JF, Nelson G, Wang C, Richter T, Simillion C, Proctor CJ, Miwa S, Olijslagers S, Hallinan J, Wipat A, Saretzki G, Rudolph KL, Kirkwood TB, von Zglinicki T. Feedback between p21 and reactive oxygen production is necessary for cell senescence. *Mol Syst Biol* 2010;6:347.

11. Warsaw G, Greber B, Falk SS, Harder C, Siatkowski M, Schordan S, Som A, Endlich N, Scholer H, Reptsilber D, Endlich K, Fuellen G. ExprEssence—revealing the essence of differential experimental data in the context of an interaction/regulation net-work. *BMC Syst Biol* 2010;4:164.
12. Xue H, Xian B, Dong D, Xia K, Zhu S, Zhang Z, Hou L, Zhang Q, Zhang Y, Han JD. A modular network model of aging. *Mol Syst Biol* 2007;3:147.
13. Kirkwood TB. Systems biology of ageing and longevity. *Phil Trans R Soc Lond Series B, Biol Sci* 2011;366:64–70.
14. Kowald A, Kirkwood TB. A network theory of ageing: the interactions of defective mitochondria, aberrant proteins, free radicals and scavengers in the ageing process. *Mutat Res* 1996;316:209–236.
15. van Leeuwen IM, Vera J, Wolkenhauer O. Dynamic energy budget approaches for modelling organismal ageing. *Phil Trans R Soc Lond Series B, Biol Sci* 2010;365:3443–3454.
16. Plank M, Wuttke D, van Dam S, Clarke SA, de Magalhaes JP. A meta-analysis of caloric restriction gene expression profiles to infer common signatures and regulatory mechanisms. *Mol BioSys* 2012;8:1339–1349.
17. Silva AS, Wood SH, van Dam S, Berres S, McArdle A, de Magalhaes JP. Gathering insights on disease etiology from gene expression profiles of healthy tissues. *Bioinformatics* 2011;27:3300–3305.
18. Li Y, de Magalhaes JP. Accelerated protein evolution analysis reveals genes and pathways associated with the evolution of mammalian longevity. *Age* 2011; doi: 10.1007/s11357-011-9361-y.
19. Vinuela A, Snoek LB, Riksen JAG, Kammenga JE. Genome-wide gene expression regulation as a function of genotype and gene in *C. elegans*. *Genome Research* 2010;20:929–937.
20. Vinuela A, Snoek LB, Riksen JA, Kammenga JE. Aging uncouples heritability and expression-QTL in *Caenorhabditis elegans*. *G3* 2012;2:597–605.
21. Langmead B, Trapnell C, Pop M, Salzberg SL. Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biol* 2009;10:R25.
22. Guthke R, Moller U, Hoffmann M, Thies F, Topfer S. Dynamic network reconstruction from gene expression data applied to immune response during bacterial infection. *Bioinformatics* 2005;21:1626–1634.
23. Langley P. An interactive environment for explanatory biological modeling, in Proceedings of the Thirty-Second Annual Meeting of the Cognitive Science Society. Portland, 2010.
24. Berger F, Ramirez-Hernandez MH, Ziegler M, The new life of a centenarian: Signalling functions of NAD(P). *Trends Biochem Sci* 2004;29:111–118.
25. Bogan KL, Evans C, Belenky P, Song P, Burant CF, Kennedy R, Brenner C. Identification of Isn1 and Sdt1 as glucose- and vitamin-regulated nicotinamide mononucleotide and nicotinic acid mononucleotide [corrected] 5'-nucleotidases responsible for production of nicotinamide riboside and nicotinic acid riboside. *J Biol Chem* 2009;284:34861–34869.
26. Schuster S, Dandekar T, Fell DA. Detection of elementary flux modes in biochemical networks: A promising tool for pathway analysis and metabolic engineering. *Trends Biotechnol* 1999;17:53–60.
27. Ruppig E, Papin JA, de Figueiredo LF, Schuster S. Metabolic reconstruction, constraint-based analysis and game theory to probe genome-scale metabolic networks. *Curr Opin Biotechnol* 2010;21:502–510.
28. de Figueiredo LF, Gossmann TI, Ziegler M, Schuster S. Pathway analysis of NAD+ metabolism. *Biochem J* 2011; 439:341–348.
29. Calin GA, Croce CM. MicroRNA signatures in human cancers. *Nat Rev Cancer* 2006;6:857–866.
30. Grillari J, Grillari-Voglauer R. Novel modulators of senescence, aging, and longevity: Small non-coding RNAs enter the stage. *Exp Gerontol* 2010;45:302–311.
31. Lai X, Schmitz U, Gupta S, Bhattacharya A, Kunz M, Wolkenhauer O, Vera J. Computational analysis of target hub gene repression regulated by multiple and cooperative miRNAs *Nucleic Acid Res* 2012;doi: 10.1093/nar/gks657.
32. Russel SJ, Kahn CR. Endocrine regulation of ageing. *Nat Rev Mol Cell Biol* 2007;8:681–691.
33. Blagosklonny MV. Molecular damage in cancer: An argument for mTOR-driven aging. *Aging* 2011;3:1130–1141.
34. Blagosklonny MV. Aging: ROS or TOR. *Cell Cycle* 2008;7: 3344–3354.
35. Holzenberger M, Dupont J, Ducos B, Leneuve P, Geloën A, Even PC, Cervera P, Le Bouc Y. IGF-1 receptor regulates lifespan and resistance to oxidative stress in mice. *Nature* 2003;421:182–187.
36. Sawitzky S, Zeissler A, Langhammer M, Biellohuby, Stock P, Hammon HM, Görs S, Metges CC, Stoehr BJM, Bidingmaier M, Fromm-Dornieden C, Baumgartner B, Christ B, Brenig B, Binder G, Metzger F, Renne U, Hoeflich A. Phenotype selection reveals coevolution of muscle glyco-gen and protein and PTEN as a gate keeper for the accretion of muscle mass in mid-aged female mice. *PLoS One* 2012;(in press).
37. Wuttke D, Connor R, C. V, Craig T, Li Y, Wood S, Vasieva O, Shmookler Reis R, Tang F, de Magalhães JP. Dissecting the gene network of dietary restriction to identify evolutionarily conserved pathways and new functional genes. *PLoS Genet* 2012;(in press).
38. Fortney K, Morgen EK, Kotlyar M, Jurisica I. In silico drug screen in mouse liver identifies candidate calorie restriction mimetics. *Rejuvenation Res* 2012;15:148–152.
39. Wang C, Maddick M, Miwa S, Jurk D, Czapiewski R, Saretzki G, Langie SA, Godschalk RW, Cameron K, von Zglinicki T. Adult-onset, short-term dietary restriction reduces cell senescence in mice. *Aging* 2010;2:555–566.
40. Bittig AT, Haack F, Maus C, Uhrmacher AM. Adapting rule-based model descriptions for simulating in continuous and hybrid space. In Proc. 9th Conf. on Computational Methods in Systems Biology. ACM, 2011, pp. 161–170.
41. Hoehndorf R, Dumontier M, Oellrich A, Wimalaratne S, Rebholz-Schuhmann D, Schofield P, Gkoutos GV, A common layer of interoperability for biomedical ontologies based on OWL EL. *Bioinformatics* 2011;27:1001–1008.

Address correspondence to:

Georg Fuellen  
 Institute for Biostatistics and Informatics in Medicine and  
 Ageing Research  
 Rembrandtstrasse 16  
 Rostock, Mecklenburg-West Pomerania, 18057  
 Germany

E-mail: fuellen@uni-rostock.de

Received: July 10, 2012

Accepted: July 11, 2012